

RESEARCH PAPER

Association Between Silver Nanoparticle Dose and Brain or Renal NF-κB Gene Expression

Gulboy A. Nasir¹, Mohammed A. Najm², Huda M. Mahmood^{3*}

¹ College of Agricultural Engineering Sciences, University of Baghdad, Iraq

² Department of Pharmacology, College of Medicine, Ibn Sina University of Medical and Pharmaceutical Sciences, Baghdad, Iraq

³ Department of Biotechnology, College of Sciences, University of Anbar, Iraq

ARTICLE INFO

Article History:

Received 05 September 2024

Accepted 27 December 2024

Published 01 January 2025

Keywords:

AgNP

CNS

NF-κB Gene

Renal abnormalities

ABSTRACT

Several studies have claimed that AgNPs' safety is not completely verified. The current work studied the CNS and renal genetic abnormalities (as manifested by modifications in NF-κB transcription) in mice treated with diverse AgNP concentrations. The findings revealed that as the amount of AgNPs was elevated, NF-κB production increased considerably.

How to cite this article

Nasir G., Najm M., Mahmood H. Association Between Silver Nanoparticle Dose and Brain or Renal NF-κB Gene Expression. J Nanostruct, 2025; 15(1):249-254. DOI: 10.22052/JNS.2025.01.024

INTRODUCTION

Nanoparticle synthesis is on the rise these days for utilization in several essential applications. [1-3]. Because of their unique features, AgNPs are broadly applied in different daily goods [4-12], but the probability of Ag⁺ ions leaking may adversely impact their safety [13-17]. As per numerous studies, the toxic effects of AgNPs and Ag⁺ release have no connection [18, 19]; nevertheless, some analysts have said that expected toxicities coming from silver nanoparticle uses may be higher than the cytotoxic effects of silver ions singly [20, 21]. NF-κB is a negative and positive regulator of gene expression [22], its activity modifies immunity, inflammatory response, and nervous system

* Corresponding Author Email: huda.mahmood@uoanbar.edu.iq

function & enhances cancer development [23, 24]. This study is carried out to see how AgNPs affected CNS and renal NF-κB expression.

MATERIAL AND METHODS

Synthesis of AgNPs

The prepared AgNPs were (40±5) nm in size and spherical form. D-sorbitol was used as a capping agent and olive leaf plant extract as a reducing agent [25, 26].

Experimental Animals

Male mus musculus (Balb/C) mice, (Fifty-six) weighting (23–35) g and aged eight weeks, were used. They came from the Al-Nahrain Centre of



This work is licensed under the Creative Commons Attribution 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

Biotechnology's animal house. They were held with a 12-hour light/dark cycle [27], at (23-25) °C, and a 7-day acclimatization period before the experiment began [28], and were randomly distributed into 7 groups, each with eighth mice in isolated cages, and provided a standard pellets meal and adequate water every day as follows:

-The first and fourth groups were given an intraperitoneal quantity of 0.25 mg/kg of AgNPs (50 μ l) for one and two weeks.

-Second, and fifth groups were given 0.5 mg/kg of silver nanoparticle solution (50 μ l). Intraperitoneal for one and two weeks, respectively.

-The third and sixth groups received a dose of 1 mg/kg body weight of AgNPs.

- The seventh group received an intraperitoneal dosage of distilled water (50 μ l) instead of an AgNP solution.

Tissue Sampling

The fourth, fifth, and sixth groups were

sacrificed the day after the treatment period ended (after 14 days), while the first, second, and third groups were sacrificed via cervical dislocation (Euthanasia) the day after the dosing period ended (after 7 days).

Tissue preparation for RNA extraction

Tissues were processed for histological evaluation using Junqueira and Carneiro's (2003) procedure, with paraffin slices used for livers, spleens, kidneys, and brains [29]. 5-8 semi-thin slices (0.5-10 μ m) were prepared from tissue paraffin blocks using an electrical ultra-microtome. The sections were used for RNA extraction.

Pre-Extraction Preparations

1- The lyophilized enzyme was dissolved in 550 μ l of RNase-free water to make the DNase I Stock Solution.

2- In the preparation of RPE buffer, absolute ethanol, 44 μ l (96-100%) was added to a container

Table 1. PCR Condition Program.

Temperature	Heated Lid	Step 1	Step 2	Step 3	Step 4	Step 5
	111 °C	25 °C	37 °C	42 °C	75 °C	4 °C
	--	10	10	60	5	∞

Table 2. Control gene (B-ACTIN) & the target gene (NF- κ B) primers.

Primers	(5 $\rightarrow$ 3)	Melting Temperature
Beta-ACTIN - F	CCTGAACCTAAGGCCAAC	60 °C
Beta-ACTIN - R	ACGTACATGGCTGGGGTGT	62 °C
NF- κ B - F	GTCAACAGATGGCCATACC	62.0 °C
NF- κ B - R	GATGGGCCTTCACACACATA	60.0 °C

Table 3. NF- κ B measured concentrations.

Period	AgNPs (mg)	Mean $\pm$ SE	
		CNS levels	Renal levels
Week	0.025 mg	0.269 $\pm$ 0.043 bc	0.081 $\pm$ 0.014 a
	0.5 mg	0.217 $\pm$ 0.066 bcd	0.069 $\pm$ 0.014 ab
	1 mg	0.296 $\pm$ 0.081 b	0.078 $\pm$ 0.001 a
Two weeks	0.025 mg	0.6128 $\pm$ 0.058 a	0.0347 $\pm$ 0.012 cd
	0.5 mg	0.1117 $\pm$ 0.017 cd	0.0164 $\pm$ 0.004 d
	1 mg	0.2982 $\pm$ 0.056 b	0.0450 $\pm$ 0.004 bc
	Control	0.0678 $\pm$ 0.064 d	0.0154 $\pm$ 0.002 d
	LSD value	0.169 **	0.027 **

** (P<0.01).

with 11 μ l of buffer RPE concentration.

3-The Qiagen Kit was used to purify total RNA from Formalin Fixed Paraffin Embedded (FFPE) tissue slices and extract RNA, which were then stored at -20°C.

cDNA Synthesis

Using an Applied Biosystem RNA-to-cDNA kit (Part No. 4387406), RNA was reverse-transcribed to complementary DNA (cDNA). For use in real-time PCR, cDNAs were stored at -20°C. RT-PCR.

Each sample was quantified in duplicate using the SYBR Green master mix. In addition, each run contained two non-template controls. Control gene mRNA quantities were employed to normalize NF- κ B mRNA levels, ensure proper cDNA synthesis,

and describe computations. Primer 3 Plus was used to generate primers for the NF- κ B gene, with B-ACTIN as a reference gene.

The change in the target's expression in a sample as compared to the same sample at time zero [30] was determined using the relative quantitation (RQ) technique, which relies on cycle threshold (CT).

A 20 μ l reaction was applied for the SYBR Green assay. A 96-well plate was given to each gene. The qRT-PCR procedure was as follows.:

Stage 1 was 50 °C / two minutes, Stage 2 was 95 °C / 10 minutes, and Stage 3 had a three-step cycle process (denaturation 95 °C for ten seconds, annealing 62 °C /60 sec., and extension 72 °C for forty-five seconds) that was performed fifty times




Fig. 1. Effect of study groups in Brain/NFKB gene.




Fig. 2. Effect of study groups in Kidney/NFKB gene.

before being cooled to 40 °C for ten seconds.

QRT-PCR Data Analysis

The threshold cycle (CT) was used to measure the fold change and gene expression levels. The reference gene (B-ACTIN gene) was used to standardize the CT data.

Δ CT sample = Ct Sample - Ct endogenous control

Δ CT calibrator = Ct control - Ct endogenous control

The relative fold change of gene expression is measured using the normalized Δ CT data and particular calibrators:

$\Delta\Delta$ CT = Δ CT sample - Δ CT calibrator

Relative copy number = Fold change = $2^{-\Delta\Delta$ CT

RESULTS AND DISCUSSION

The study found varying doses and durations of AgNPs administration affected tissue expression levels of the NF- κ B gene in many organs, including the brain and kidneys (Table 3).

The lowest value of NF- κ B gene expression in the brain was 0.1117, which was obtained from the lowest AgNPs dose (0.025 mg/kg) for two weeks ($P < 0.01$), while the highest value was (0.6128) only after daily administration of the lowest AgNPs dose (0.025 mg/kg of body weight) for two weeks ($P < 0.01$), as shown in Fig. 1.

After one week of treatment with the highest dose of AgNPs (0.025 mg/kg of body weight), kidney NF- κ B gene expression was 0.081 ($P < 0.01$), but after two weeks at a dose of 0.5 mg/kg, kidney expression was 0.164 ($P < 0.01$). as shown in Fig. 2.

Utilizing products even if they contain a small amount of silver, can have a bad impact on immunity and general health. [31-35]. The recent findings discovered that variation in the given AgNP doses led to different elevated CNS levels of NF- κ B, regardless of the duration of mice exposure.

Prolog exposure to the excitatory glutamatergic stimuli was demonstrated to produce a harmful NF- κ B activation in the neurons [36-41].

It has been found that the non-regular NF- κ B activation or abnormal production of NF- κ B can lead to many CNS abnormalities, like modification of the apoptotic response to neurodegenerative cells and, an enhance the inflammation of neurons

that may lead to neuronal cell death [42].

The current investigation found that the varied supplied AgNP dosages resulted in varying concentrations of renal NF- κ B expression during the two periods of mice exposure.

The NF- κ B contribution to the incidence of acute kidney injury or other kidney damage has been recognized over time [43–51]. First of all, NF- κ B affects the renal expression of inflammatory genes [52, 53], furthermore, cytokines, genotoxic stress, mechanical stress, mechanical stress, and other mediators are among the triggers that activate NF- κ B in kidney injury [54, 55].

The tumor suppressor protein, p53, interrupts the cell cycle and promotes cell death in response to DNA damage and cellular stress [56, 57]. A current study found that the lack of p53 during kidney damage confers a protective effect [58–61]. Increasing evidence suggests that p53 has a κ B site, implying that NF- κ B can bind to it and control p53 gene production [62, 63]. Which in turn, enhances the inflammatory and apoptotic response which leads to further kidney damage.

CONCLUSION

The results here revealed a substantial relationship between NF- κ B expression and AgNP administered dose, suggesting a potential risk associated with utilizing AgNP goods.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interests regarding the publication of this manuscript.

REFERENCES

1. Al-taei EH. Effect of Silver Nanoparticles Synthesized Using Leaves Extract of Olive on Histopathological and Cytogenetic Effects in Albino Mice. *Iraqi Journal of Agricultural Sciences*. 2020;51(5):1448-1457.
2. Nowack B, Bucheli TD. Occurrence, behavior and effects of nanoparticles in the environment. *Environ Pollut*. 2007;150(1):5-22.
3. Sun TY, Mitrano DM, Bornhöft NA, Scheringer M, Hungerbühler K, Nowack B. Envisioning Nano Release Dynamics in a Changing World: Using Dynamic Probabilistic Modeling to Assess Future Environmental Emissions of Engineered Nanomaterials. *Environmental Science and Technology*. 2017;51(5):2854-2863.
4. Hussein K. Detection of the Antimicrobial Activity of Silver Nanoparticles Biosynthesized By *Streptococcus Pyogenes* Bacteria. *Iraqi Journal of Agricultural Sciences*. 2020;51(2):500-507.
5. Durán N, Durán M, de Jesus MB, Seabra AB, Fávaro WJ, Nakazato G. Silver nanoparticles: A new view on mechanistic

- aspects on antimicrobial activity. *Nanomed Nanotechnol Biol Med*. 2016;12(3):789-799.
6. Kumar SSD, Houreld NN, Kroukamp EM, Abrahamse H. Cellular imaging and bactericidal mechanism of green-synthesized silver nanoparticles against human pathogenic bacteria. *J Photochem Photobiol B: Biol*. 2018;178:259-269.
7. Tian X, Jiang X, Welch C, Croley TR, Wong T-Y, Chen C, et al. Bactericidal Effects of Silver Nanoparticles on Lactobacilli and the Underlying Mechanism. *ACS Applied Materials and Interfaces*. 2018;10(10):8443-8450.
8. Capek I. Preparation of metal nanoparticles in water-in-oil (w/o) microemulsions. *Advances in Colloid and Interface Science*. 2004;110(1-2):49-74.
9. Sohn EK, Johari SA, Kim TG, Kim JK, Kim E, Lee JH, et al. Aquatic Toxicity Comparison of Silver Nanoparticles and Silver Nanowires. *BioMed research international*. 2015;2015:893049-893049.
10. Silver Nanoparticle Applications. *Engineering Materials: Springer International Publishing*; 2015.
11. Wei L, Lu J, Xu H, Patel A, Chen Z-S, Chen G. Silver nanoparticles: synthesis, properties, and therapeutic applications. *Drug Discovery Today*. 2015;20(5):595-601.
12. McGillicuddy E, Murray I, Kavanagh S, Morrison L, Fogarty A, Cormican M, et al. Silver nanoparticles in the environment: Sources, detection and ecotoxicology. *Sci Total Environ*. 2017;575:231-246.
13. Ratte HT. Bioaccumulation and Toxicity of Silver Compounds: A Review. *Environmental Toxicology and Chemistry*. 1999;18(1):89.
14. Yue Y, Li X, Sigg L, Suter MJF, Pillai S, Behra R, et al. Interaction of silver nanoparticles with algae and fish cells: a side by side comparison. *Journal of nanobiotechnology*. 2017;15(1):16-16.
15. Li L, Wu H, Ji C, van Gestel CAM, Allen HE, Peijnenburg WJGM. A metabolomic study on the responses of daphnia magna exposed to silver nitrate and coated silver nanoparticles. *Ecotoxicology and Environmental Safety*. 2015;119:66-73.
16. Sakamoto M, Ha J-Y, Yoneshima S, Kataoka C, Tatsuta H, Kashiwada S. Free Silver Ion as the Main Cause of Acute and Chronic Toxicity of Silver Nanoparticles to Cladocerans. *Archives of Environmental Contamination and Toxicology*. 2014;68(3):500-509.
17. Shen M-H, Zhou X-X, Yang X-Y, Chao J-B, Liu R, Liu J-F. Exposure medium: key in identifying free Ag⁺ as the exclusive species of silver nanoparticles with acute toxicity to *Daphnia magna*. *Sci Rep*. 2015;5:9674-9674.
18. Fabrega J, Luoma SN, Tyler CR, Galloway TS, Lead JR. Silver nanoparticles: Behaviour and effects in the aquatic environment. *Environ Int*. 2011;37(2):517-531.
19. Sørensen SN, Hjorth R, Delgado CG, Hartmann NB, Baun A. Nanoparticle ecotoxicity—physical and/or chemical effects? Integrated Environmental Assessment and Management. 2015;11(4):722-724.
20. Pakrashi S, Tan C, Wang W-X. Bioaccumulation-based silver nanoparticle toxicity in *Daphnia magna* and maternal impacts. *Environmental Toxicology and Chemistry*. 2017;36(12):3359-3366.
21. Abramenko NB, Demidova TB, Abkhalimov EV, Ershov BG, Krysanov EY, Kustov LM. Ecotoxicity of different-shaped silver nanoparticles: Case of zebrafish embryos. *J Hazard Mater*. 2018;347:89-94.
22. Mattson MP, Culmsee C, Yu Z, Camandola S. Roles of Nuclear Factor κ B in Neuronal Survival and Plasticity. *J Neurochem*. 2000;74(2):443-456.
23. Mahmood AS, Farhan SH, Najm MA. Association of Polymorphism GST1 Gene and Antioxidant status, and Interleukin-17 of Colorectal Cancer Iraqi Patients. *Indian Journal of Forensic Medicine and Toxicology*. 2019;13(4):535.
24. Albensi BC. What Is Nuclear Factor Kappa B (NF- κ B) Doing in and to the Mitochondrion? *Frontiers in cell and developmental biology*. 2019;7:154-154.
25. Gulboy A. Nasir, Alaa K. Mohammed, Hasan F. Samir, 2016. Biosynthesis and Characterization of Silver Nanoparticles Using Olive Leaves Extract and Sorbitol. *Iraqi Journal of Biotechnology*, 15, (1), 22-32.
26. Nasir, G.A., Najm, M.A., Hussein, A.L., 2020. The effect of silver nanoparticles on braf gene expression. *Systematic Reviews in Pharmacy*, 11(10), pp. 570–575.
27. Wolfensohn S. *Laboratory Animals: An Introduction for Experimenters*, 2nd edition Edited by A A Tuffery (1995). John Wiley and Sons Ltd: Chichester. 392 pp. Hardback. Obtainable from the publishers, Baffins Lane, Chichester, West Sussex PO19 1UD, UK (ISBN 0 471 95257 5). Price £39.95. *Anim Welfare*. 1996;5(1):85-85.
28. Mostafa RM, Mostafa YM, Ennaceur A. Effects of exposure to extremely low-frequency magnetic field of 2 G intensity on memory and corticosterone level in rats. *Physiology and Behavior*. 2002;76(4-5):589-595.
29. McGurk S. *Junqueira's Basic Histology Text and Atlas – 13th edition* Mescher Anthony L Junqueira's Basic Histology Text and Atlas – 13th edition 544pp + CD-ROM £45.99 McGraw-Hill Professional 978 1 2590 7232 1 1259072320. *Nurs Stand*. 2013;28(16):34-34.
30. Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2- $\Delta\Delta$ CT Method. *Methods*. 2001;25(4):402-408.
31. Pratap Singh S, Sharma S. Arsenic exposure and its toxicity. *Current Trends in Pharmacy and Pharmaceutical Chemistry*. 2022;4(1):24-29.
32. Najm, M.A., Alfatlawi, W.R., Saleh, R.O., Kadhim Al Taie, S.M., 2019. Evaluation of antioxidants (GSH, CAT, SOD) and MDA in Iraqi women with breast cancer. *Journal of Global Pharma Technology*, 11(7), pp. 107–110.
33. Nasir GA, Khudhair IA, Najm MA, Mahmood HM. Nanotechnology at the Molecular Level. *Al-Rafidain Journal of Medical Sciences (ISSN: 2789-3219)*. 2022;3:71-74.
34. I.A., Al-Essawi, Iman Abbas, H.M., Mahmood, Huda M. Effect of Gold Nanoparticles on hmgA Gene Expression of *Pseudomonas aeruginosa* Isolates. *Journal of Nanostructures*, 2024.
35. Ghaidaa Raheem Lateef Al-Awsi, Gulboy Nasir, Majida M. Meteab Alshammari, Abdolmajid Ghasemian. (2020). Antibacterial and Antibiofilm Effects of Bismuth Nanoparticles Produced by *Bacillus Subtilis* Against Multidrug-Resistant *Pseudomonas Aeruginosa*. *Malaysian Journal of Biochemistry and Molecular Biology* 3:15-22.
36. Kaltschmidt C, Kaltschmidt B, Baeruerle PA. Stimulation of ionotropic glutamate receptors activates transcription factor NF-kappa B in primary neurons. *Proceedings of the National Academy of Sciences of the United States of America*. 1995;92(21):9618-9622.
37. Guerrini L, Blasi F, Denis-Donini S. Synaptic activation of NF-kappa B by glutamate in cerebellar granule neurons in vitro. *Proceedings of the National Academy of Sciences of the United States of America*. 1995;92(20):9077-9081.

38. Detect the Infection with Rubella Virus and Toxoplasmosis in Pregnancy Causes Suffering from Early Abortion by Using Real Time PCR. Medico-Legal Update. 2020.
39. Caccamo D, Campisi A, Marini H, Adamo EB, Li Volti G, Squadrito F, et al. Glutamate promotes NF- κ B pathway in primary astrocytes: protective effects of IRFI 016, a synthetic vitamin E analogue. *Exp Neurol*. 2005;193(2):377-383.
40. Sitcheran R, Comb WC, Cogswell PC, Baldwin AS. Essential role for epidermal growth factor receptor in glutamate receptor signaling to NF- κ B. *Molecular and cellular biology*. 2008;28(16):5061-5070.
41. Correlating Schizophrenia with DRD3 Ser9Gly or HTR2 Receptor Gene Variants by using RFLP Method. *Indian Journal of Forensic Medicine and Toxicology*. 2020.
42. Dresselhaus EC, Meffert MK. Cellular Specificity of NF- κ B Function in the Nervous System. *Front Immunol*. 2019;10:1043-1043.
43. Akcay A, Nguyen Q, Edelstein CL. Mediators of inflammation in acute kidney injury. *Mediators Inflamm*. 2009;2009:137072-137072.
44. Tadagavadi RK, Reeves WB. Renal dendritic cells ameliorate nephrotoxic acute kidney injury. *Journal of the American Society of Nephrology : JASN*. 2010;21(1):53-63.
45. Linkermann A, De Zen F, Weinberg J, Kunzendorf U, Krautwald S. Programmed necrosis in acute kidney injury. *Nephrology Dialysis Transplantation*. 2012;27(9):3412-3419.
46. López-Franco O, Suzuki Y, Sanjuán G, Blanco J, Hernández-Vargas P, Yo Y, et al. Nuclear factor-kappa B inhibitors as potential novel anti-inflammatory agents for the treatment of immune glomerulonephritis. *The American journal of pathology*. 2002;161(4):1497-1505.
47. Francescato HDC, Costa RS, Scavone C, Coimbra TM. Parthenolide reduces cisplatin-induced renal damage. *Toxicology*. 2007;230(1):64-75.
48. Hazim, O.H., Najm, M.A., Ibraheem, Z.O., 2020. Genetic diversity of three medicinal plants of the apiaceous family. *Systematic Reviews in Pharmacy*, 11(8), pp. 775–778.
49. Ramesh G, Reeves WB. p38 MAP kinase inhibition ameliorates cisplatin nephrotoxicity in mice. *American Journal of Physiology-Renal Physiology*. 2005;289(1):F166-F174.
50. Sanz AB, Sanchez-Niño MD, Ramos AM, Moreno JA, Santamaria B, Ruiz-Ortega M, et al. NF- κ B in Renal Inflammation. *J Am Soc Nephrol*. 2010;21(8):1254-1262.
51. Cao CC, Ding XQ, Ou ZL, Liu CF, Li P, Wang L, et al. In vivo transfection of NF- κ B decoy oligodeoxynucleotides attenuate renal ischemia/reperfusion injury in rats. *Kidney Int*. 2004;65(3):834-845.
52. Plümpe J, Malek NP, Bock CT, Rakemann T, Manns MP, Trautwein C. NF- κ B determines between apoptosis and proliferation in hepatocytes during liver regeneration. *American Journal of Physiology-Gastrointestinal and Liver Physiology*. 2000;278(1):G173-G183.
53. Sanz AB, Sanchez-Niño MD, Ramos AM, Moreno JA, Santamaria B, et al., 2010. NF- κ B in renal inflammation. *J Am Soc Nephrol*, 21: 1254–1262.
54. Guijarro C, Egido J. Transcription factor- κ B (NF- κ B) and renal disease. *Kidney Int*. 2001;59(2):415-424.
55. Chen G, Shaw MH, Kim Y-G, Nuñez G. NOD-Like Receptors: Role in Innate Immunity and Inflammatory Disease. *Annual Review of Pathology: Mechanisms of Disease*. 2009;4(1):365-398.
56. Wei Q, Dong G, Yang T, Megyesi J, Price PM, Dong Z. Activation and involvement of p53 in cisplatin-induced nephrotoxicity. *American journal of physiology Renal physiology*. 2007;293(4):F1282-F1291.
57. Zhou L, Fu P, Huang XR, Liu F, Lai KN, Lan HY. Activation of p53 promotes renal injury in acute aristolochic acid nephropathy. *Journal of the American Society of Nephrology : JASN*. 2010;21(1):31-41.
58. Gudkov AV, Gurova KV, Komarova EA. Inflammation and p53: A Tale of Two Stresses. *Genes Cancer*. 2011;2(4):503-516.
59. Sutton TA, Hato T, Mai E, Yoshimoto M, Kuehl S, Anderson M, et al. p53 is renoprotective after ischemic kidney injury by reducing inflammation. *Journal of the American Society of Nephrology : JASN*. 2013;24(1):113-124.
60. Daghdan B, Kaddar N, Ayyed Najm M, Barakat H. Comparison of double dose of Paracetamol tablets 500mg with one Paracetamol tablet 1000mg. *Research Journal of Pharmacy and Technology*. 2024;4671-4676.
61. Kim G, Ouzounova M, Quraishi AA, Davis A, Tawakkol N, Clouthier SG, et al. SOCS3-mediated regulation of inflammatory cytokines in PTEN and p53 inactivated triple negative breast cancer model. *Oncogene*. 2015;34(6):671-680.
62. Wu H, Lozano G. NF- κ B activation of p53. A potential mechanism for suppressing cell growth in response to stress. *J Biol Chem*. 1994;269(31):20067-20074.
63. Jeong S-J, Radonovich M, Brady JN, Pise-Masison CA. HTLV-I Tax induces a novel interaction between p65/RelA and p53 that results in inhibition of p53 transcriptional activity. *Blood*. 2004;104(5):1490-1497.